

Roll No.

Total Pages : 04

GSO/M-20
MATHEMATICS
BM-362
Linear Algebra

1722

Time : Three Hours]

[Maximum Marks : 26

Note : Attempt *Five* questions in all, selecting *one* question from each Section. Q. No. **1** is compulsory.

Compulsory Question

1. (a) What can you say about the linear span of the empty set ? 1
- (b) Let $T : U \rightarrow V$ be a homomorphism, then prove that $\ker T$ is a subspace of U . 1
- (c) Show that $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ given by $T(x, y) = (2x - y, x - y, -2x)$ is a linear transformation. 1
- (d) Normalize the vector $u = (2, 1, -1)$ in \mathbb{R}^3 . 1
- (e) Show that the linear transformation $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ defined by $T(x, y, z) = (x + z, x - z, y)$ is non-singular. 2

Section I

2. (a) Prove that a minimal generating set of a finitely generated vector space $V(F)$ is always a basis of V . $2\frac{1}{2}$
- (b) Show that the union $W_1 \cup W_2$ of subspaces of a vector space V need not be a subspace of V . $2\frac{1}{2}$
3. (a) Let $U = L(S_1)$ and $V = L(S_2)$, where :
 $S_1 = \{(1, 3, -2, 2, 3), (1, 4, -3, 4, 2), (2, 3, -1, -2, 9)\}$,
 $S_2 = \{(1, 3, 0, 2, 1), (1, 5, -6, 6, 3), (2, 5, 3, 2, 1)\}$.
Find basis and dimension of $U + V$. $2\frac{1}{2}$
- (b) Let W be a subspace of a finite dimensional vector space $V(F)$, then show that $\dim \frac{V}{W} = \dim V - \dim W$. $2\frac{1}{2}$

Section II

4. (a) Find a linear transformation $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ such that $T(1, 1, 1) = (1, 0)$ and $T(1, 1, 2) = (1, -1)$. $2\frac{1}{2}$
- (b) If $T : U \rightarrow V$ be a linear transformation, then show that $\dim(R(T)) + \dim(N(T)) = \dim U$. $2\frac{1}{2}$
5. (a) Find a linear transformation $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ whose range is generated by $(1, 0, -1), (1, 2, 2)$. $2\frac{1}{2}$

- (b) If $B = \{(-1, 1, 1), (1, -1, 1), (1, 1, -1)\}$ be a basis of \mathbb{R}^3 , then find the dual basis of B . **2½**

Section III

6. (a) Let $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be a linear operator defined by $T(x, y, z) = (2x, 4x - y, 2x + 3y - z)$. Show that T is invertible and find T^{-1} . **2½**
- (b) Find the linear map $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ whose matrix is

$$A = \begin{bmatrix} 1 & -1 \\ -2 & 3 \\ 0 & 1 \end{bmatrix}, \text{ relative to the ordered basis}$$

- $B = \{(1, 1), (0, 2)\}$ and $B' = \{(0, 1, 1), (1, 0, 1), (1, 1, 0)\}$ for \mathbb{R}^3 . **2½**
7. (a) Find the co-ordinates of $(1, 2, 1)$ relative to the basis $\{(1, 1, 2), (2, 2, 1), (1, 2, 2)\}$ using change of basis matrix (transition matrix). **2½**
- (b) If T be an invertible operator and λ is an eigen value of T , then show that λ^{-1} is an eigen value of T^{-1} . **2½**

Section IV

8. (a) Let V be an inner product space, then show that $\|u + v\| = \|u\| + \|v\|$. **2½**

- (b) Obtain an orthonormal basis with respect to standard inner product for the subspace of \mathbb{R}^3 generated by $(1, 0, 1)$, $(1, 0, -1)$ and $(0, 3, 4)$. $2\frac{1}{2}$
9. (a) Show that a linear operator T on a unitary space V is Hermitian iff $\langle T(\alpha), \alpha \rangle$ is real for every α . $2\frac{1}{2}$
- (b) Let T be a linear operator on an inner product space $V(F)$. If $T^2(u) = 0$ and T is self-adjoint or skew-symmetric, then show that $T(u) = 0$. $2\frac{1}{2}$